

研究実績一覧

英文

1. Horita Y, Tadokoro M, Taura K, Ashida R, Hiu M, Taguchi T, Furusu A, Kohno. Prednisolone co-administered with losartan confers renoprotection in patients with IgA nephropathy. *Renal Failure* 2007;29:441-446.
2. Horita Y, Taura K, Taguchi T, Furusu A, Kohno. Aldosterone breakthrough during therapy with angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers in proteinuric patients with IgA nephropathy. *Nephrology (Carlton)* 2006;11:462-466.
3. Horita Y, Yakabe K, Tadokoro M, et al. Renal circulatory effects of acetazolamide in patients with essential hypertension. *Am J Hypertens* 2006;19:282-285.
4. Horita Y, Tadokoro M, Taura K, Suyama N, Taguchi T, Miyazaki M, Kohno S. Low dose-combination therapy with temocapril and losartan reduces proteinuria in normotensive patients with immunoglobulin A nephropathy. *Hypertens Res* 2004;27:963-970.
5. Horita Y, Tadokoro M, Taura K, Suyama N, Taguchi T, Miyazaki M, Kohno S. Histologically confirmed superimposition of post-streptococcal acute glomerulonephritis during IgA nephropathy. *Clin Exp Nephrol* 2004;8:351-355.
6. Horita Y, Tadokoro M, Taura K, Suyama N, Nazneen A, Taguchi T, Miyazaki M, Kohno S. Incidental detection of renomedullary interstitial cell tumour in a renal biopsy specimen. *Nephrol Dial Transplant* 2004;19:1007-1008.
7. Horita Y, Nakamura S, Hirashima S, Tadokoro M, Taura K, Sakamoto I, Nishikido M, Miyazaki M, Kohno S. Evaluation of Tc-99m MAG3 dynamic renal scintigraphy with an aneurysmal type renal arteriovenous fistula. *Clin Nucl Med* 2003;28:429-431.
8. Horita Y, Hayashida K, Fukuchi K, Takishita S, Tadokoro M, Taura K, Suyama N, Miyazaki M, Kohno S, Kawano Y. Acetazolamide assisted Tc-99m MAG3 renography to assess renal blood flow reserve. *Ann Nucl Med* 2003;17:139-144.
9. Horita Y, Tadokoro M, Taura S, Suyama N, Miyazaki M, Taguchi T, Harada T, Ozono Y, Kohno S. Renal outcome of immunoglobulin A nephropathy with mild proteinuria. *Acta Med Nagasaki* 2003;48:39-43.
10. Horita Y, Tadokoro M, Suyama N, Taura K. Severity of glomerulosclerosis predicts prognosis of IgA nephropathy with proteinuria. *Acta Med Nagasaki* 2002;47:123-126.
11. Horita Y, Tadokoro M, Taura K, Mishima Y, Miyazaki M, Kohno S, Kawano Y. Relationship between Carotid Artery Intima-Media Thickness and Atherosclerotic Renal Artery Stenosis in Type 2 Diabetes with Hypertension. *Kidney Blood Press Res* 2002;25:255-259.
12. Horita Y, Tsunoda S, Inenaga T, Kawano Y, Ishibashi-Ueda H, Chiba Y, Takishita S. Pregnancy outcome in nephrotic syndrome with mixed connective tissue disease. *Nephron* 2001;89:354-356.

13. Horita Y, Hayashida K, Takishita S, Kohno S, Kawano Y. Dilatation of renal artery distal to stenosis demonstrated using acetazolamide Tc-99m MAG3 scintigraphy. *Clin Nucl Med* 2001;26:795-796.
14. Horita Y, Noguchi M, Miyazaki M, Tadokoro M, Taura K, Watanabe T, Nishiura K, Harada T, Ozono Y, Kohno S. Prognosis of patients with rounded atelectasis undergoing long-term hemodialysis. *Nephron* 2001;88:87-92.
15. Horita Y, Inenaga T, Nakahama H, Ishibashi-Ueda H, Kawano Y, Nakamura S, Horio T, Okuda N, Ando M, Takishita S. Cause of residual hypertension after adrenalectomy in patients with primary aldosteronism. *Am J Kidney Dis* 2001;37:884-889.
16. Nakahama H, Sakaguchi K, Horita Y, Sasaki O, Nakamura S, Inenaga T, Takishita S. Treatment of severe hypothyroidism reduced serum creatinine levels in two chronic renal failure patients. *Nephron* 2001;88:264-267.
17. Fukuchi K, Inenaga T, Suzuki Y, Horita Y, Hayashida K, Ishida Y. Paraganglioma seen with FDG Dual –Head Gamma Camera coincidence imaging after false-negative results of I-123 MIBG imaging. *Clin Nucl Med* 2001;26:966-967.
18. Horita Y, Miyazaki M, Kadota J, Watanabe T, Yamashita M, Nishiura K, Taguchi T, Matsuo T, Ozono Y, Kohno S. Type II diabetes mellitus and primary Sjögren's syndrome complicated by pleural effusion. *Intern Med* 2000;39:979-984.
19. Horita Y, Miyazaki M, Noguchi M, Tadokoro M, Taura K, Ozono Y, Kohno S. Healing of Fournier's gangrene of the scrotum in a haemodialysis patient after conservative therapy alone. *Nephrol Dial Transplant* 2000;15:419-421.
20. Horita Y, Miyazaki M, Koji T, Kobayashi N, Shibuya M, Razzaque MS, Cheng M, Ozono Y, Kohno S, Taguchi T. Expression of vascular endothelial growth factor and its receptors in rats with protein-overload nephrosis. *Nephrol Dial Transplant* 1998;13:2519-2528.
21. Horita Y. Ultrastructural study of the peritoneum in patients on continuous ambulatory peritoneal dialysis. *Acta Med Nagasaki* 1997;42:5-11.
22. Tsuda N, Chowdhury PR, Hayashi T, Anami M, Iseki M, Koga S, Matsuya F, Kanetake H, Saito Y, Horita Y. Primary renal angiosarcoma: a case report and review. *Pathol Int* 1997;47:778-783.
23. Ikematsu Y, Usui K, Kamohara Y, Inoue K, Maeda J, Amano M, Horita Y. Leiomyoma of the lesser omentum: report of a case. *Surg Today* 1996;26:46-48.
24. Razzaque MS, Cheng M, Horita Y, Nishihara M, Harada T, Taguchi T. Immunohistochemical analysis of Type III and IV collagens in tubulointerstitial damage in human benign nephrosclerosis. *J Int Med Res* 1995;23:480-486.
25. Razzaque MS, Koji T, Horita Y, Nishihara M, Harada T, Nakane PK, Taguchi T. Synthesis of type III collagen and type IV collagen by tubular epithelial cells in diabetic nephropathy. *Pathol Res Pract* 1995;191:1099-1104.

和文

1. 堀田義雄、田所正人、田浦幸一、中村 貴、塚崎祥子、宮崎雅也、須山尚史、平島 定、宮崎正信、河野 茂. 急性膵炎により高血糖性意識障害発症した維持血液透析患者の 1 例. 透析会誌 2003;36:1273-1277.
2. 堀田義雄、Nazneen A、程 明、Razzaque MS、浪江 智、田所正人、田浦幸一、原田孝司、田口 尚. 微小変化型ネフローゼ症候群を合併した全身性エリテマトーデスの 1 症例. 日腎誌 1997;39:759-764.